



# Lake Crenshaw

LAKE HABITAT ASSESSMENT, LAKE VEGETATION INDEX, SUBMERGED  
VEGETATION SURVEY AND WATER QUALITY

David Eilers, Ray Dominguez | USF Water Institute | May 21, 2020

# Methods

## STUDY AREA ANALYSIS

The watershed containing the Lake Crenshaw was analyzed using ESRI ArcGIS 10.6. Using this software with 2017 Hillsborough County aerial, 2014 Land Use/ Land Cover (LULC), Landscape Development Intensity (LDI) Index values were calculated for the 100 meter buffer surrounding the lake following the procedures of Reiss & Brown 2012 (Reiss & Brown, 2012. Landscape Development Intensity (LDI) Index User's Manual. H.T. Odum Center for Wetlands, University of Florida. March 2012). According to Reiss and Brown "The LDI represents a human disturbance gradient for wetland systems. The LDI is an integrated measure of human activity, combining the effects from air and water pollutants, physical damage, changes in the suite of environmental conditions ... on the structure and processes of landscapes and ecosystems... Natural, undeveloped LU/LC classes have a LDI index value of one. In the Florida framework, the maximum LDI index score is approximately 42."

The LDI is calculated by multiplying each land use coefficient by the percentage of the area of interest occupied by that land use, and then summing the results. The Florida Department of Environmental Protection (DEP) uses the LDI as a tool to estimate potential land use impacts on streams, lakes, and wetlands. LDI values less than two ( $\leq 2$ ) can be considered minimally disturbed."

## LAKE BATHYMETRY AND MORPHOLOGICAL CHARACTERISTICS ASSESSMENT

The **Bathymetric Map**<sup>1</sup> provides the lake's morphologic parameters in various units. The bottom of the lake was mapped using a Lowrance Elite 7 Ti Wide Area Augmentation System (WAAS)<sup>2</sup> enabled Global Positioning System (GPS) with Totalscan transducer (bottom sounder) to determine the boat's position, and bottom depth in a single measurement. The result is an estimate of the lake's area, mean and maximum depths, and volume and the creation of a bottom contour map. Besides pointing out the deeper fishing holes in the lake, the morphologic data derived from this part of the assessment can be valuable to overall management of the lake vegetation as well as providing flood storage data for flood models.

---

<sup>1</sup> A bathymetric map is a map that accurately depicts all of the various depths of a water body. An accurate bathymetric map is important for effective herbicide application and can be an important tool when deciding which form of management is most appropriate for a water body. Lake volumes, hydraulic retention time and carrying capacity are important parts of lake management that require the use of a bathymetric map.

<sup>2</sup> WAAS is a form of differential GPS (DGPS) where data from 25 ground reference stations located in the United States receive GPS signals from GPS satellites in view and retransmit these data to a master control site and then to geostationary satellites. For more information, see end note 2.

## LAKE VEGETATION INDEX ASSESSMENT

Hillsborough County requested the implementation of the Florida Department of Environmental Protection methods for Lake Vegetation Index (LVI 1000) (<http://www.dep.state.fl.us/water/sas/sop/sops.htm>) using forms FD 9000-03 (Physical/Chemical Characterization), FD 9000-06 (Lake Habitat Assessment) FD 9000-27 (LVI Field Sheet) and FD 9000-31 (Lake Observation Field Sheet).

The Lake Vegetation Index (LVI) is a rapid assessment protocol in which selected sections of a lake are assessed for the presence or absence of vegetation through visual observation and through the use of a submerged vegetation sampling tool called a Frodus. The assessment results provide a list of species presents and the dominant and where appropriate co-dominant species that are found in each segment. These results are then entered into a scoring table and a final LVI score is determined. LVI scores provide an estimate of the vegetative health of a lake. Our assessment team was trained and qualified by FDEP to conduct these assessment as an independent team and must prequalify each year prior to conducting additional assessments. The LVI method consists of dividing the lake into twelve pie-shaped segments (see diagram below) and selecting a set of four segments from the twelve to include in the LVI. The assessment team then travels across the segment and identifies all unique species of aquatic plant present in the segment. Additionally, a Frodus is thrown at several points on a single five-meter belt transect that is established in the center of the segment from a point along the shore to a point beyond the submerged vegetation zone. For scoring, the threshold score for impairment is 43.

Four metrics are utilized in the Lake Vegetation Index Survey; Dominant Coefficient of Conservatism (CoC), Percent Florida Exotic Pest Plant Council Type 1 (% FLEPPC), Percent Native Taxa, Percent Sensitive Taxa.

The Dominant Coefficient of Conservatism (CoC) metric for the dominant or co-dominate species in each section. The CoC applies a score of 0-10 to each species based on its ecological tolerances and fidelity to pre-settlement conditions. Species with higher scores show a high fidelity to native, undisturbed habitats and are typically sensitive to alterations. Available CoC scores can be obtained from LT 7000 from the Florida Department of Environmental Protection at: <http://www.dep.state.fl.us/water/sas/sop/sops.htm>.

The percent FLEPPC (Florida Exotic Pest Plant Council) Category 1 invasive exotic taxa in a single sampling unit (pie slice) by dividing the number of FLEPPC Category I taxa by the total number of taxa in that sampling unit. Multiply result times 100. Refer to Appendix LVI 1000-1 to determine which plants are on the FLEPPC Category 1 list. Note that not all exotic taxa should be included in this metric, only those listed in Appendix LVI 1000-1 as Category 1 FLEPPC. If the FLEPPC updates their list of Category 1 exotics, those updates shall not be reflected in this calculation until they are included in Appendix LVI 1000-1.

The percent native taxa in a single sampling unit (pie slice) is calculated by dividing the number of native taxa by the total number of taxa in that sampling unit. Multiply result times 100. Nativity status is determined by the Plant Atlas from the Institute for Systematic Botany, and is listed in Appendix LVI 1000-1. For informational purposes, visit the website

<http://www.florida.plantatlas.usf.edu/>. Taxa that are native according to the Plant Atlas from the Institute for Systematic Botany but are not on the list in Appendix LVI 1000-1 may be included in this metric calculation, but inclusion of these additional taxa is not required.

The percent sensitive taxa in a single sampling unit by summing the number of taxa with a C of C (Coefficient of Conservatism) score  $\geq 7$  and then dividing by the total number of taxa in that sampling unit. Multiply result times 100. Refer to Appendix LVI 1000-1 for a list of C of C scores.

The collected bathymetric data is analyzed for submerged aquatic vegetation (SAV) calculations including the percentage of the surface area of the lake inhabited by SAV as well as an estimate of the percent volume of the lake inhabited by SAV. SAV is an important component to a lakes nutrient cycling as well as chlorophyll concentrations due to the SAV and phytoplankton competing for available nutrients in the water column. In addition SAV serves a vital role as habitat for many species of macroinvertebrates and fish as well as substrate for epiphytic algae.

## WATER QUALITY ASSESSMENT


Physical water quality samples were taken using a Eureka Manta Sub-2 multiprobe pre and post calibrated on the day of the assessment. Measurements taken with this device include: depth, conductivity, pH, Dissolved Oxygen (mg/l and % Saturation) and salinity. Chemical water parameters were collected and preserved on ice by USF Water Institute staff and analyzed at the Environmental Protection Commission of Hillsborough County Laboratory. Analysis include; Chlorophyll (a, b, c, t and corrected), Alkalinity, Color, EColi, Enterococci, Ammonia, Nitrates/Nitrites, Total Phosphorous, Kjeldahl Nitrogen and Total Nitrogen. The results of the water quality sampling effort will be discussed in the framework of the FDEP Numeric Nutrient Criteria



## Study Area

Lake Crenshaw is located in north-western Hillsborough County, Florida. The Landscape Development Intensity Index of the 100 meter buffer around Lake Crenshaw is dominated by Residential (61.8%) and Natural (34.4%) land uses. The resulting LDI value for the 100 meter buffer around Lake Crenshaw is 5.18.

*FIGURE 1: 2020 Lake Crenshaw ASSESSMENT STUDY AREA MAP*



## Lake Bathymetry and Morphological Characterization

At the time of the assessment, Lake Crenshaw was experiencing elevated water levels (53.70 feet above sea level NAVD 88) resulting in a 31.2 acre water body. Lake Crenshaw at the time of the assessment had a mean water depth of 8.6 feet and a maximum observed depth of 15.84 feet. The volume at this time was approximately 88,006,533 gallons. Figure 2 shows the resulting bathymetric contour map for Lake Crenshaw from data collected on May 21, 2020. The collected data has been overlain the 2020 Hillsborough County aerals.

*Table 1: Morphological Calculations for Lake Crenshaw*

Parameter	Feet	Meters	Acres	Acre-Ft	Gallons
Surface Area (sq)	1,360,328	126,378	31.2		
Mean Depth	8.6	2.64			
Maximum Depth	15.84	4.83			
Volume (cubic)	11,764,681	333,136		270.1	88,006,533
Gauge (NAVD 88)	53.7	16.37			

Figure 2: 2020 1-Foot Bathymetric Contour Map for Lake Crenshaw

## Lake Crenshaw

### Depth (feet)

15 - 16  
14 - 15  
13 - 14  
12 - 13  
11 - 12  
10 - 11  
9 - 10

8 - 9  
7 - 8  
6 - 7  
5 - 6  
4 - 5  
3 - 4  
2 - 3  
1 - 2  
0 - 1

### EXPLANATION:

Survey Date: May 21, 2020  
Water level was 53.70 feet above sea level NAVD 88 at the time of the assessment.  
Contours are expressed in absolute depth below this level.

### LAKE MORPHOLOGY:

Perimeter 4,670 ft;  
Area 31.2 Acres;  
(1,360,328 Sqft)  
Mean Depth 8.6 ft;  
Volume 270.1 Acre-ft,  
(88,006,533 gallons);  
Deepest point 15.84 ft

### DATA SOURCES:

2020 aerial photography provided by ESRI.  
Lake perimeter digitized from Hillsborough County 2020 aerial photographs.  
All contours generated by the USF Water Institute from survey data collected by USF Water Institute Lake and Stream

### DISCLAIMER:

This map is for illustrative purposes only, and should not be used for lake navigation.

0 100 200 400  
Feet






Figure 3 Overview photograph of Lake Crenshaw showing typical shoreline vegetation in an undeveloped area.



## Lake Habitat and Lake Vegetation Index Assessment

The lake assessment for Lake Crenshaw was conducted on May 21, 2020. The water in Lake Crenshaw was characterized as moderately tannic and moderately turbid. The secchi disk depth was 2.4 feet in part due to the high tannins and turbidity. The vegetation quality of the plants in and buffering Lake Chapman are predominantly native species with moderate growths of non-native nuisance species such as *Ludwigia peruviana*, *Panicum repens* and *Melaleuca quinquenervia*. Most stormwater reaches the lake via sheet flow. The bottom substrate quality was dominated by sand with coarse particulate organic matter near shore and some accumulation of muck. Approximately 61.8% of the surrounding land has been developed for residential housing including several docks and seawalls. Some homeowners have maintained a vegetated buffer zone along the shoreline.






*Figure 4 Cephalanthus occidentalis, Common Buttonbush, along a vegetated section of Lake Crenshaw shoreline.*

The Lake Vegetation Index identified 34 species of wetland vegetation growing in the four selected sections along Lake Crenshaw. The majority of these species (26) are native species. The remaining 8 species (*Panicum repens*, *Cyperus prolifer*, *Melaleuca quinquenervia*, *Schinus terebinthifolius*, *Sphagneticola trilobata*, *Colocasia esculenta*, *Sapium sebiferum*, and *Ludwigia peruviana*) are non- native and invasive to this region. The vegetation community along Lake Crenshaw is dominated by a variety of emergent species including *Panicum repens*, *Acer rubrum*, *Panicum hemitomon*, *Melaleuca quinquenervia* and *Cephalanthus occidentalis*. The water's surface in Lake Crenshaw was dominated by *Nuphar*. A total of 4 species of submerged aquatic vegetation were observed, *Utricularia gibba*, *Chara*, *Nitella* and *Potamogeton diversifolius* with *Chara* being the dominant species. Submerged vegetation was limited in the lake due to the low water visibility and tanins blocking available light. By analyzing the collected sonar chart, submerged aquatic vegetation potentially covered approximately 2.67% of the surface area of Lake Crenshaw. This submerged vegetation inhabits an estimated 0.05% of the water volume in Lake Crenshaw. Figure 5 shows the results of the SAV analysis indicating the location and height of SAV canopy.

The calculated LVI score for Lake Crenshaw was 43, equal to the impairment threshold of 43 indicating that the vegetation community is "Healthy". Figure 6 shows the map of Lake Crenshaw detailing the LVI regions used for the assessment (Regions 2, 5, 8, 11). Table 2 details the species list results of the Lake Vegetation Index. Table 3 details the scoring result for the Lake Vegetation Index.

# Lake Crenshaw

## Height of Submerged Vegetation



Lake Perimeter  
Ground Level

## EXPLANATION:

Survey Date: May 21, 2020  
Water level was 53.70' NAVD 88 at the time of the assessment. Submerged Aquatic Vegetation was analyzed from collected sonar data. The height of the SAV where present is shown in 0.25 foot increments.

## DATA SOURCES:

2020 aerial photography provided by ESRI.  
Lake perimeter digitized from Hillsborough County 2020 aerial photographs. All contours generated by the USF Water Institute from survey data collected by USF Water Institute Lake and Stream Assessment Program.

## SAV STATISTICS:

Area 36,292 square ft; 0.8 Acres;  
(2.67% of Lake Surface Area)  
Mean SAV Height 0.2 ft;  
Volume 6,246 Cubic ft, (46,726 gallons);  
(0.05 % of Lakes Volume)



Water Institute



0 125 250 500 Feet



Figure 5 Lake Crenshaw Submerged Aquatic Vegetation Assessment Results






Figure 6: Lake Vegetation Index region map for Lake Crenshaw

Table 2: Lake Vegetation Index results for Lake Crenshaw May 21, 2020

Species	CofC	Region			
		2	5	8	11
<i>Acer rubrum</i>	4.65	1	1	1	1
<i>Blechnum serrulatum</i>	5.5	1	1	1	1
<i>Cephalanthus occidentalis</i>	5	1	1	1	1
<i>Chara</i>	3.9	D	1	C	D
<b><i>Melaleuca quinquenervia</i></b>	0	1	1	1	1
<i>Mikania scandens</i>	1.95	1	1	1	1
<i>Nuphar</i>	3.5	1	1	1	1
<i>Panicum hemitomon</i>	5.82	1	D	C	1
<b><i>Panicum repens</i></b>	0	1	1	1	1
<i>Pontederia cordata</i>	5.38	1	1	1	1
<i>Sagittaria lancifolia</i>	3	1	1	1	1
<i>Taxodium</i>	7	1	1	1	1
<b><i>Ludwigia peruviana</i></b>	0	1	1		1
<i>Utricularia gibba</i>	6.37	1	1		1
<b><i>Colocasia esculenta</i></b>	0		1		1
<i>Nitella</i>	6	1		1	
<i>Canna flaccida</i>	5.5		1		
<i>Centella asiatica</i>	1.92			1	
<b><i>Cyperus prolifer</i></b>	0		1		
<i>Diodia virginiana</i>	3			1	
<i>Hydrocotyle</i>	2			1	
<i>Ilex cassine</i>	6		1		
<i>Myrica cerifera</i>	2				1
<i>Najas guadalupensis</i>	5.07			1	
<i>Nymphaea odorata</i>	5				1
<i>Persea palustris</i>	7			1	
<i>Pluchea baccharis</i>	5.45				1
<i>Potamogeton diversifolius</i>	6			1	
<i>Quercus laurifolia</i>	4			1	
<b><i>Sapium sebiferum</i></b>	0	1			
<b><i>Schinus terebinthifolius</i></b>	0				1
<b><i>Sphagneticola trilobata</i></b>	0			1	
<i>Triadenum virginicum</i>	5		1		
<i>Typha</i>	1	1			

Table 3: Scoring Summary for the Lake Vegetation Index

LVI Score Summary	Region			
	2	5	8	11
Total # of taxa in sampling unit	17	19	21	19
% Native taxa in sampling unit	76.47059	73.68421	85.71429	73.68421
% FLEPPC CAT 1 taxa in sampling unit	23.52941	21.05263	9.52381	26.31579
% Sensitive taxa in sample unit	5.882353	5.263158	9.52381	5.263158
Dominant CoC in sample unit	3.9	5.82	4.86	3.9
Native Score $((x-62.5)/37.5)$ or $((x-66.67)/25.89)=$	0.378547	0.270924	0.735585	0.270924
Invasive FLEPPC 1 Score $(1 - (x/30))=$	0.215686	0.298246	0.68254	0.122807
Sensitive Score $(x/(27.78 \text{ or } 20)) =$	0.294118	0.263158	0.47619	0.263158
Dominant CoC Score $(x/(7.91 \text{ or } 7)) =$	0.557143	0.831429	0.694286	0.557143
Raw Score Total = N+I+S+D =	1.445494	1.663756	2.5886	1.214031
Division Factor = (3 D=0 or 4) =	4	4	4	4
Average LVI dividend = Raw /DF	0.361374	0.415939	0.64715	0.303508
South				
LVI Score for sampling unit =	36.13735	41.59389	64.71501	30.35078
Total LVI SCORE =	43			



## Water Quality Assessment

Limited long-term water quality data is available for Lake Crenshaw. The available data was collected by USGS, Hillsborough County and University of Florida LAKEWATCH program (1968-2020), however this dataset is incomplete with recent years having sparse data. There were no samples available for 2018 and 2019. Four sampling events occurred in 2020. Table 4 provides a summary of the Physical/Chemical conditions recorded at the middle of the Lake Crenshaw.

*Table 4: Lake Crenshaw Water Quality (Field)*

Depth (m)	Temp °C	pH	DO (mg/L)	DO (%sat)	Cond (unho/cm)	Salinity (ppt)	Secchi Depth (m)
0.17	31.23	7.44	8.21	109.7	87.5	0.04	1.80
1.9	29.7	7.13	7.83	101.9	87	0.04	
3.62	26.3	6.29	0	0	92.1	0.04	
POR	24.4	6.34	7.63	98.4	55.7	0.04	1.75

The chemical water quality analysis for Lake Crenshaw is shown in Table 5 for the sample taken on June 1, 2020. Table 6 includes this data in the numeric nutrient criteria framework using the data from this assessment as well as the available LAKEWATCH geometric mean values for the period of record since complete data for the past three years for available parameters is not available. Total Phosphorous values were below the nutrient threshold for colored lakes in the west central region with insufficient data developed by FDEP of 0.05 mg/l with a value of 0.017 mg/l for the POR and equal to the threshold for the most recent samples in 2020 with a value of 0.050 mg/l. If sampling were to be sufficient (previous three years of quarterly sampling) the threshold could be as high as 0.16 mg/L.

Total Nitrogen values were below the nutrient threshold for colored lakes with insufficient data developed by FDEP of 1.27 mg/l with a value of 0.676 mg/l for the POR data. The Total Nitrogen value for the most recent 2020 data was 0.540 mg/l. If sampling were to be sufficient (previous three years of quarterly sampling) the threshold could be as high as 2.23 mg/L. Chlorophyll-a corrected values are below the nutrient threshold for colored lakes developed by FDEP of 20.0 µg/l with a value of 2.4 µg/l, however this data is only available for the 4 samples taken in 2020. The UF LAKEWATCH data is for uncorrected chlorophyll-a and has a mean value for the period of record of 7.5 µg/l.

2020 bacteria testing showed low levels of E. Coli (10.5 colonies/100ml) and Enterococci (21.1 colonies/100ml) below the rules set forth in FDEP 62-302.530 (<https://www.flrules.org/gateway/RuleNo.asp?title=SURFACE%20WATER%20QUALITY%20STANDARDS&ID=62-302.500>) “Most Probable Number (MPN) or Membrane Filter (MF) counts shall

not exceed a monthly average of 200, nor exceed 400 in 10% of the samples, nor exceed 800 on any one day. Monthly averages shall be expressed as geometric means based on a minimum of 10 samples taken over a 30 day period.”

*Table 5: Lake Crenshaw Water Quality Results from 6/1/20 (Laboratory)*

Parameter	Lake Crenshaw (Center)	POR Mean Value	Units
Alkalinity		3.52	mg/LCaCO <sub>3</sub>
E Coli	5.2		#/100ml
Nitrates/Nitrites	0.043	0.034	mg/L
Enterococci	2	21.2	#/100 ml
Chlorophyll a	4.2	7.50	ug/L
Chlorophyll b	1.1	1.1	ug/L
Chlorophyll c	1.3	1.1	ug/L
Chlorophyll t			ug/L
Chlorophylla Corr	2.5	2.4	ug/L
Chlorophyll-pheo			ug/L
Ammonia	0.046	0.063	mg/L
Kjeldahl Nitrogen	0.560	0.767	mg/L
Total Nitrogen	0.560	0.676	mg/L
Total Phosphorus	0.050	0.017	mg/L
Color(345)F.45		52.7	Pt/Co

Table 6: Numeric Nutrient Criteria Framework

Parameter	Value
Geometric Mean (Geomean) Color (pcu)	52.7
Number of Samples	23
Geometric Mean Alkalinity (mg/L CaCO <sub>3</sub> )	3.53
Number of Samples	7
Lake Type	Colored
Chlorophyll a Criteria (ug/L)	20
Insufficient for Geomean Criteria then P mg/L	0.05
Insufficient for Geomean Criteria then N mg/L	1.27
Geomean Chla Corrected ug/L	2.4
Geomean TP mg/L	0.017
Geomean TN mg/L	0.676
Number of Samples	428 (POR)
Potential Impaired Chlorophyll a	Not Impaired
Potential Impaired TP	Not Impaired
Potential Impaired TN	Not Impaired



## Conclusion

The results of the assessment of Lake Crenshaw shows a healthy lake based on Total Nitrogen and Total Phosphorous concentrations according to the FDEP numeric nutrient criteria using the 2020 samples taken during this assessment and limited long term water quality record. The sampling data was insufficient to calculate proper FDEP Numeric Nutrient Criteria values. Consistent Long term sampling would be necessary to determine actual NNC values with a minimum of three samples per year for the previous three years. The most recent year of data for Total Phosphorous is below the nutrient threshold. The system also shows partial health in the vegetation communities according to the Lake Vegetation Index with moderate overall species (34), moderate occurrences of non-native, invasive species and few sensitive plant species with an overall LVI score of 43. The assessment also revealed some submerged aquatic vegetation community comprising 4 species occupying 2.67% of the surface area and 0.05% of the volume of Lake Crenshaw.