



Hollomans Branch

STREAM HABITAT ASSESSMENT, STREAM CONDITIONS INDEX, LINEAR
VEGETATION SURVEY, RAPID PERIPHYTON SURVEY AND WATER QUALITY

David Eilers, Megan Dunne, Lauren Kier | USF Water Institute | April 21, 2021

Methods

STUDY AREA ANALYSIS

The watershed containing the stream being assessed was analyzed using ESRI ArcGIS 10.2. Using this software with 2020 Hillsborough County aerial, 2017 Land Use/ Land Cover (LULC) and Waterbody ID (WBID) layers courtesy of the Florida Department of Environmental Protection (FDEP). The Landscape Development Intensity Index (LDI) was calculated for the WBID containing the stream. From FDEP

(<https://floridadep.gov/dear/bioassessment/content/bioassessment-ldi-hdg-bcg>) “The Landscape Development Intensity index (LDI) is an estimate of how much humans have altered an area of interest around a waterbody. Various land use types (low density residential, row crops, industrial and natural) are assigned coefficients of land use intensity based on estimates of the amount of human energy that is put into those land use types.”

The LDI is calculated by multiplying each land use coefficient by the percentage of the area of interest occupied by that land use, and then summing the results. The FDEP uses the LDI as a tool to estimate potential land use impacts on streams, lakes, and wetlands. LDI values less than two (≤ 2) can be considered minimally disturbed.” In the Florida framework, the maximum LDI index score is approximately 42.

HABITAT AND VEGETATION ASSESSMENT

For small streams that are not easily navigated by Jonboat for bathymetric mapping and vegetation analysis, Hillsborough County requested the implementation of the FDEP methods for Stream and River Habitat Assessment (FT 3100)

(<http://www.dep.state.fl.us/water/sas/sop/sops.htm>) using forms FD 9000-3, FD 9000-4 and FD 9000-5, Rapid Periphyton Survey (FS 7230) using form FD 9000-25 and Linear Stream Vegetation Survey (FS 7320) using form FD 9000-32. These methods were utilized on two sampling locations on each stream, typically near access points along roadways.

Stream and River Habitat Assessment per FT3100 receives a score calculated in Form FD 9000-5. This score results from the ranking of the primary habitat components (substrate diversity, substrate availability, water velocity and habitat smothering) and secondary habitat components (Artificial channelization, bank stability, riparian buffer zone width and riparian zone vegetation quality). The maximum score possible in this method is a 160.

Two metrics are utilized in the Linear Vegetation Survey (LVS). The Mean Coefficient of Conservatism (CoC) applies a score of 0-10 to each species based on its ecological tolerances and fidelity to pre-settlement conditions. Species with higher scores show a high fidelity to native, undisturbed habitats and are typically sensitive to alterations. Available CoC scores can be obtained from Table LVI 1000-1 from the Florida Department of Environmental Protection at: <http://www.dep.state.fl.us/water/sas/sop/sops.htm>. The Percent Florida Exotic Pest Plant Council (% FLEPPC) metric calculates the percent invasive exotics as the number of occurrences of FLEPPC Category I or II in the 100 m reach divided by the total number of taxa occurrences in the 100 m reach. The FLEPPC list can be found in FDEP LVI 1000-1.

STREAM CONDITION INDEX ASSESSMENT

The Stream Condition Index (SCI) was sampled and calculated per DEP SOP SCI 1000. . The SCI consists of collecting macroinvertebrates via 20 D-frame dipnet sweeps (0.5 m in length) in the most productive habitats in a 100 m reach of stream. The organisms are sub-sampled, and identified to the lowest practical taxonomic level. The SCI is composed of ten metrics, eight of which decrease in response to human disturbance, with two metrics (% very tolerant and % dominant) increasing in response to human disturbance. According to DEP SOP SCI 1000, the SCI scores greater than 35 are considered healthy. Proposed biological health assessment criteria state that a WBID is considered to meet designated uses if the average of the two most recent SCI scores is 40 or higher and neither of the most recent of those scores is less than 35.

WATER QUALITY ASSESSMENT

Physical water quality samples were taken using a Eureka Manta Sub-2 multiprobe pre and post calibrated daily. Measurements taken with this device include: depth, conductivity, pH, Dissolved Oxygen (mg/l and % Saturation) and salinity. Chemical water parameters were collected and preserved on ice by USF Water Institute staff and analyzed at the Hillsborough County Public Utilities Laboratory. Analysis include; Chlorophyll (a, b, c, t and corrected), Alkalinity, Color, E. Coli, Enterococci, Ammonia, Nitrates/Nitrites, Total Phosphorous, Kjeldahl Nitrogen and Total Nitrogen. Results will be discussed in the Florida Department of Environmental Protection's Numeric Nutrient Criteria framework and combined with the monthly sampling from the Hillsborough County Environmental Protection Commission Monthly sampling data.

Study Area

Hollomans Branch is located in north eastern Hillsborough County in the Hillsborough Bay Watershed. Its headwaters are located south of Sam Allen and De Montmollin Road. The outfall of Hollomans Branch is in Flint Creek. The assessment of Hollomans Branch was conducted on April 21, 2021 downstream of the US Highway 301 Bridge. At the time of the assessment, the water levels were normal for the dry season. The Hollomans Branch WBID covers 22.0 square miles and is dominated by residential (39.7%), agricultural (30.6%) and natural (23.1%) land uses. The resulting calculated landscape development intensity index score was 4.37.






Figure 2 Overview photograph of the Hollomans Branch Sample Site showing the typical habitat features

Habitat and Vegetation Assessment

The region of Hollomans Branch where the assessment was conducted is in a natural region downstream from the US Highway 301 Bridge. The region was heavily shaded with a mean canopy cover measurement of 92.4%. Hollomans Branch averaged 0.26 meters in depth, approximately 3.47 meters wide with a flow of 0.32 m/s.

The primary habitat components of the FDEP Habitat Assessment focus on in-water habitat. The primary habitat components score in the optimal category for Water Velocity (0.32 m/s). Suboptimal category scores were achieved for Habitat Smothering (sufficient pools but many of the productive habitats were affected by sand smothering). Substrate Diversity (Presence of two major productive habitats (snags, roots)) was scored as marginal. Substrate Availability (3.05% of stream are productive habitats) was scored as poor. Minor habitats included leaf packs/mats, rocks and sand and silt deposits. The total score for the primary habitat components was a 43 out of 80.

The secondary habitat components of the FDEP Habitat Assessment focus on the surrounding features of the stream. The secondary habitat components scored in the optimal category for Bank Stability (both banks with few raw eroded areas and surrounded by expansive floodplain forest), Artificial Channelization, Riparian Zone Vegetation Quality (both banks showing low levels of disturbance shown in the species present) and Riparian Buffer Zone Width (greater than 18 meters of buffer). The secondary habitat components received a score of 76 out of 80. The resulting FDEP Habitat Assessment score was a 119.

Table 1 Scoring Summary for the Stream Habitat Assessment

Metric		Score
Primary Habitat Components		
	Substrate Diversity	10
	Substrate Availability	3
	Water Velocity	19
	Habitat Smothering	11
	Primary Score	43
Secondary Habitat Components		
	Artificial Channelization	20
	Bank Stability - Right Bank	9
	Bank Stability - Left Bank	9
	Riparian Buffer Zone Width - Right Bank	10
	Riparian Buffer Zone Width - Left Bank	10
	Riparian Zone Vegetation Quality - Right Bank	9
	Riparian Zone Vegetation Quality - Left Bank	9
	Secondary Score	76
Habitat Assessment Score		119

Periphyton was not encountered during the 99 samples taken during the Rapid Periphyton Survey. The tree canopy in the assessment area averaged 92.4% reducing available light for periphyton to flourish.

The FDEP Linear Vegetation Survey encountered less than 2 m² of herbaceous species rooted in Hollomans Branch at the time of the assessment. The vegetation surrounding the creek was dominated by *Ulmus* and *Quercus*.

Table 1 Linear Vegetation Survey Results – Hollomans Branch

Taxa Name	C of C Score	Sample Site										Total Occurrences
		0-10	10-20	20-30	30-40	40-50	50-60	60-70	70-80	80-90	90-100	
<i>Less than 2m²</i>												



Figure 3 Snags were the most abundant major productive habitat in Hollomans Branch.

Stream Condition Index

The analysis of the SCI sample involves splitting the sample into 2 aliquots for analysis. The SCI metrics are then calculated on each separately. The final SCI score is an average of the two scores. The SCI score for Hollomans Branch was 58 out of a possible 100 points, corresponding with a “Category 2 Healthy” designation, with minor loss of taxonomic diversity from the expected community of a healthy stream. Both subsamples contained moderate total taxa with 22 taxa in both subsample A and B. The most recent previous SCI sample for Hollomans Branch at this sample site was collected on 12/14/2017 with a score of 72.

High scores (scores above 7.0) were achieved for the % Dominance and % Tanytarsini in both subsamples. Additional high scores were achieved for Total Trichoptera in Sample B and % Very Tolerant Individuals in Sample A. Low scores (less than 3.0) were achieved for the Total Taxa in both subsamples. In Sample A, Total Trichoptera and Total Sensitive Taxa also scored low. In Sample B, Total long-lived Taxa scored low. The full results of the SCI sampling are shown in Table 3 (Sample A) and Table 4 (Sample B) for Hollomans Branch.

Table 2 SCI metric summaries for Hollomans Branch Sample A (top) and Sample B (bottom)

SCI Metric	Raw Totals	SCI scores	Adjusted SCI scores
Total Taxa	22.00	2.92	2.92
Total Ephemeroptera	2.00	4.00	4.00
Total Trichoptera	2.00	2.86	2.86
% Filter Feeders	28.39	6.44	6.44
Total Clingers	3.00	4.29	4.29
Total Long-lived Taxa	1.00	3.33	3.33
% Dominance	19.35	8.93	8.93
% Tanytarsini	15.48	8.24	8.24
Total Sensitive Taxa	1.00	1.43	1.43
% Very Tolerant Individuals	4.52	7.48	7.48

SCI Sum	49.91
Final SCI score	55.46

SCI Metric	Raw Totals	SCI scores	Adjusted SCI scores
Total Taxa	22.00	2.92	2.92
Total Ephemeroptera	2.00	4.00	4.00
Total Trichoptera	5.00	7.14	7.14
% Filter Feeders	27.48	6.23	6.23
Total Clingers	4.00	5.71	5.71
Total Long-lived Taxa	0.00	0.00	0.00
% Dominance	19.87	8.83	8.83
% Tanytarsini	17.88	8.64	8.64
Total Sensitive Taxa	3.00	4.29	4.29
% Very Tolerant Individuals	5.96	6.90	6.90

SCI Sum	54.66
Final SCI score	60.73

Table 3 SCI full results for Sample A

Stream Condition Index Results for Holloman Branch SCIA																				
Phylum	Subphylum	Class	Subclass	Order	Family	Taxa	Abundance	Collapsed Abundance	Taxa Presence	Ephemeroptera Taxa	Trichoptera Taxa	50% Filterer	100% Filterer	Clinger Taxa	Long-lived Taxa	Dominant Taxa	Tanytarsini	Sensitive Taxa	Very Tolerant Individuals	Specimen Notes
Mollusca		Gastropoda	Heterobranchia	Hygrophila	Ancylidae	<i>Ferrissia fragilis</i>	2	2	1	0	0	0	0	0	0	0	0	0	0	
Mollusca		Gastropoda	Heterobranchia	Hygrophila	Physidae	<i>Physa acuta</i>	1	1	1	0	0	0	0	0	0	0	0	0	1	
Mollusca		Bivalvia	Autobranchia	Venerida	Cyrenidae	<i>Corbicula</i> spp.	1	1	1	0	0	0	0	1	0	1	0	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Ephemeroptera	Caenidae	<i>Caenis</i> spp.	2		0	0	0	0	0	0	0	0	0	0	0	No hind legs; not C. hilaris; 1 = damaged
Arthropoda	Hexapoda	Insecta	Pterygota	Ephemeroptera	Caenidae	<i>Caenis diminuta</i>	9	11	1	1	0	0	0	0	0	0	0	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Ephemeroptera	Baetidae	<i>Baetis intercalaris</i>	2	2	1	1	0	0	0	0	0	0	0	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Odonata	Zygoptera		1		0	0	0	0	0	0	0	0	0	0	0	Head with no PM, like Coenagrionidae
Arthropoda	Hexapoda	Insecta	Pterygota	Odonata	Coenagrionidae	<i>Argio</i> spp.	1		0	0	0	0	0	0	0	0	0	0	0	No caudal lamellae, Ant. like A. sedula
Arthropoda	Hexapoda	Insecta	Pterygota	Odonata	Coenagrionidae	<i>Argio sedula</i>	1	3	1	0	0	0	0	0	0	0	0	0	3	
Arthropoda	Hexapoda	Insecta	Pterygota	Trichoptera	Leptoceridae	<i>Nectopsyche pavidia</i>	2	2	1	0	1	0	0	0	0	0	0	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Trichoptera	Hydropsychidae	<i>Cheumatopsyche</i> spp.	27	27	1	0	1	0	0	27	1	0	0	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Coleoptera	Elmidae	<i>Dubiraphia</i> spp.	13	13	1	0	0	0	0	0	0	0	0	0	0	Adults = 13
Arthropoda	Hexapoda	Insecta	Pterygota	Coleoptera	Elmidae	<i>Stenelmis</i> spp.	4	4	1	0	0	0	0	0	1	0	0	0	0	Larvae = 2, Adults = 2
Arthropoda	Hexapoda	Insecta	Pterygota	Coleoptera	Elmidae	<i>Microcylopusus</i> spp.	22	22	1	0	0	0	0	0	0	0	0	0	0	Larvae = 15, Adults = 7
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Chironomidae</i> spp.	2		0	0	0	0	0	0	0	0	0	0	0	Damaged
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Chironomus</i> spp.	1	1	1	0	0	0	0	0	0	0	0	0	0	1
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Cladotanytarsus</i> spp.	1	1	1	0	0	0.5	0	0	0	0	1	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Tanytarsus</i> spp.	22	23	1	0	0	11.5	0	0	0	0	0	23	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Polypedium flavum</i>	29	30	1	0	0	0	0	0	0	0	0	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Ablabesmyia mallochii</i>	1	1	1	0	0	0	0	0	0	0	0	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Pentaneura inconspicua</i>	1	1	1	0	0	0	0	0	0	0	0	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Stenochironomus</i> spp.	3	3	1	0	0	0	0	0	0	0	0	0	0	
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Polypedium beckae</i>	1	1	1	0	0	0	0	0	0	0	0	0	1	
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Atrichopogonidae	<i>Atrichopogon</i> spp.	1	1	1	0	0	0	0	0	0	0	0	0	0	Larva = 1
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Tipulidae	<i>Tipula</i> spp.	1	1	1	0	0	0	0	0	0	0	0	0	0	Larva = 1
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Simuliidae	<i>Simulium</i> spp.	4	4	1	0	0	0	0	4	1	0	0	1	0	Larvae = 4

Table 4 SCI full results for Sample B

Stream Condition Index Results for Holloman Branch SCIB																				
Phylum	Subphylum	Class	Subclass	Order	Family	Taxa	Abundance	Collapsed Abundance	Taxa Presence	Ephemeroptera Taxa	Trichoptera Taxa	50% Filterer	100% Filterer	Clinger Taxa	Long-lived Taxa	Dominant Taxa	Tanytarsini	Sensitive Taxa	Very Tolerant Individuals	Specimen Notes
Annelida		Citellata	Oligochaeta	Tubificidae	Naididae	<i>Nais communis</i>	1	1	1	0	0	0	0	0	0	0	0	0	0	1
Annelida		Citellata	Oligochaeta	Tubificidae	Naididae	<i>Nais pardalis</i>	1	1	1	0	0	0	0	0	0	0	0	0	0	1
Mollusca		Gastropoda	Heterobranchia	Hygrophila	Ancylidae	<i>Ferrissia fragilis</i>	2	2	1	0	0	0	0	0	0	0	0	0	0	0
Mollusca		Gastropoda	Heterobranchia	Hygrophila	Physidae	<i>Physa acuta</i>	2	2	1	0	0	0	0	0	0	0	0	0	0	2
Mollusca		Gastropoda	Heterobranchia	Hygrophila	Planorbidae	<i>Menetus dilatatus</i>	3	3	1	0	0	0	0	0	0	0	0	0	0	3
Arthropoda	Hexapoda	Insecta	Pterygota	Ephemeroptera	Caenidae	<i>Caenis diminuta</i>	5	5	1	1	0	0	0	0	0	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Ephemeroptera	Heptageniidae	<i>Heptageniidae</i> spp.	3	3	1	1	0	0	0	1	0	0	0	1	0	No 7th gill
Arthropoda	Hexapoda	Insecta	Pterygota	Odonata	Zygoptera spp.		1	1	1	0	0	0	0	0	0	0	0	0	0	Damaged
Arthropoda	Hexapoda	Insecta	Pterygota	Trichoptera	Leptoceridae	<i>Oecetis</i> sp. A	1	1	1	0	1	0	0	0	0	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Trichoptera	Leptoceridae	<i>Nectopsyche</i> spp.	1	1	1	0	1	0	0	0	0	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Trichoptera	Leptoceridae	<i>Triaenodes</i> spp.	1	1	1	0	1	0	0	0	0	0	0	1	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Trichoptera	Hydropsychidae	<i>Cheumatopsyche</i> spp.	28	28	1	0	1	0	0	28	1	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Trichoptera	Hydroptilidae	<i>Neotrichia</i> spp.	5	5	1	0	1	0	0	0	0	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Coleoptera	Elmidae	<i>Dubiraphia</i> spp.	5	5	1	0	0	0	0	0	0	0	0	0	0	Adults = 5
Arthropoda	Hexapoda	Insecta	Pterygota	Coleoptera	Elmidae	<i>Stenelmis</i> spp.	3	3	1	0	0	0	0	0	1	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Coleoptera	Elmidae	<i>Microcyloepus</i> spp.	28	28	1	0	0	0	0	0	0	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Diptera spp.		3		0	0	0	0	0	0	0	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Chironomidae</i> spp.	3		0	0	0	0	0	0	0	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Tanytarsus</i> spp.	25	27	1	0	0	0	13.5	0	0	0	0	27	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Cryptochironomus</i> spp.	1	1	1	0	0	0	0	0	0	0	0	0	0	1
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Polypedium flavum</i>	26	30	1	0	0	0	0	0	0	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Stenochironomus</i> spp.	1	1	1	0	0	0	0	0	0	0	0	0	0	0
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Chironomidae	<i>Polypedium beckae</i>	1	1	1	0	0	0	0	0	0	0	0	0	0	1
Arthropoda	Hexapoda	Insecta	Pterygota	Diptera	Empididae	<i>Hemerodromia</i> spp.	1	1	1	0	0	0	0	0	0	0	0	1	0	Larva = 1

Water Quality Assessment

Long-term water quality data is not available for Hollomans Branch. The data that is available was collected by the FDEP (1976-2005), Hillsborough County Environmental Protection Commission (1976-2009) and Hillsborough County Public Works (2021). Values for the physical water parameters begin in 1976 and continue through 2021. Values for the laboratory water parameters begin in 1976 through 2021. The 2021 USF Water Institute Assessment fall within the range of the previous data collections. Table 5 provides a summary of the Physical/Chemical conditions recorded at the site.

Table 5 Hollomans Branch Physical Water Quality (Field)

Hollomans Branch								
Date	Depth (m)	Temp (°C)	pH	DO (mg/L)	DO (% Sat)	Cond (UMHO/cm)	Salinity (PPT)	Secchi Depth (m)
4/21/2021	0.6	22.85	6.82	7.08	81.4	211.2	0.1	0.5
Mean POR	0.5	21.49	7.06	7.03	76.87	211.1	0.24	0.26

The chemical water quality analysis for Hollomans Branch is shown in Table 6 along with mean values for the period of record for available parameters. The previous 3-years of data is incomplete as only 4 samples were taken (all in 2021). The 2021 geometric mean values for Total Phosphorous values were below the nutrient region threshold developed by FDEP of 0.49 mg/L with a geometric mean value of 0.290 mg/L (2021). Total Phosphorous values for the sample from this assessment were 0.351 mg/L. Total Nitrogen values were below the nutrient region threshold developed by FDEP of 1.65 mg/L for the 2021 data with a mean value of 1.210 mg/L (2021). The Total Nitrogen value from the assessment was below the threshold with a concentration of 1.410 mg/L. Chlorophyll-a corrected values fall below the site specific evaluation range of 3.2 µg/l to 20 µg/l for the 2021 data at 1.0 µg/l. For sites with Chlorophyll-a values in this range, the assessment is indicating conditions reflecting an balance in flora.

An elevated biomass of the bacterial parameters was observed in the 2021 dataset with E. Coli having a geomean of 531 colonies/100 ml, 1,656/100 ml for Enterococci.

Table 6 Hollomans Branch Water Quality (Laboratory)

Parameter	Hollomans Branch 4/21/2021	POR Mean (1976- 2021)	Units
Alkalinity	52.0	57.5	mg/LCaCO ₃
Color(345)F.45	60	91.0	Pt/Co
E. Coli	944	531.3	#/100 ml
Enterococci	1,840	4,589	#/100 ml
Chlorophyll a	6.1	6.97	ug/L
Chlorophyll b	1.0	1.57	ug/L
Chlorophyll c	1.0	2.28	ug/L
Chlorophyll t	6.10	3.86	ug/L
Chlorophylla Corr	1.0	1.97	ug/L
Chlorophyll-pheo	10.1	6.669	ug/L
Ammonia	< 0.073	0.037	mg/L
Kjeldahl Nitrogen	1.210	0.519	mg/L
Total Nitrogen	1.410	1.703	mg/L
Nitrates/Nitrites	0.196	0.399	mg/L
Total Phosphorus	0.351	0.315	mg/L

Conclusion

Hollomans Branch at US Highway 301 is located in a predominantly natural area. At the time of the habitat assessment, the water levels were low, corresponding to the middle of the dry season, however sufficient habitat for macroinvertebrates was observed. Due to these factors, the Habit Assessment resulted in a suboptimal score of 119. Disruption to the vegetation community was not observed in the results of the Linear Vegetation Survey with Hollomans Branch having below 2 square meters of rooted herbaceous vegetation. Hollomans Branch did meet standards for the rapid periphyton survey with 0% of samples being ranked between 4 and 6 due to the heavy canopy coverage in the region. The recent water quality record for Hollomans Branch showed concentrations of Chlorophyll-a corrected, Total Phosphorous and Total Nitrogen below the FDEP thresholds. The results of the SCI sampling indicate that the stream is “Healthy” based on the macroinvertebrate community. Elevated bacterial biomass was observed in the most recent and long term sampling. Table 7 summarizes the results of the nutrient sampling, floristic sampling, habitat assessment and SCI.

Table 7 Summary of Water Quality, Floristic Surveys and Habitat Assessments

Measure		Hollomans Branch	2019	2020	2021	Threshold
Total Phosphorous (mg/l)		0.351	N/A	N/A	0.290	< 0.49
Total Nitrogen (mg/l)		1.410	N/A	N/A	1.210	< 1.65
RPS (% Rank 4-6)		0%				< 25%
LVS	Avg C of C	N/A				≥ 2.5
	FLEPPC %	N/A				< 25%
Chlorophyll-a Corrected (µg/l)		1.0	N/A	N/A	1.0	< 20 µg/l
Habitat Assessment		119				> 34
SCI		58				> 34