

Stream Assessment Report for Delaney Creek in Hillsborough County, Florida

Date Assessed: June 30, 2014

Assessed by: David Eilers, Camille Courant, Jay Gilbert, Matt Jarrett

INTRODUCTION

This assessment was conducted to update existing physical and ecological data for Delaney Creek on the [Hillsborough County Water Atlas](#). The project is a collaborative effort between the University of South Florida's Water Institute and Hillsborough County Stormwater Management Section. The project is funded by Hillsborough County. The project has, as its primary goal, the rapid assessing of up to 150 lakes and streams in Hillsborough County during a five-year period. The product of these investigations will provide the County, property owners and the general public a better understanding of the general health of Hillsborough County lakes and streams, in terms of shoreline development, water quality, morphology (bottom contour, volume, area, etc.) and the plant biomass and species diversity. These data are intended to assist the County and its citizens to better manage lakes and streams.



Figure 1. Delaney Creek downstream from 36th Avenue South.

BACKGROUND

This study of Delaney Creek focuses on the region from the mouth in Hillsborough bay to S 86th Street. This covers the tidal portion of Delaney Creek as well freshwater portions. Delaney Creek has been altered and straightened in this study area resulting in steep banks and straight channels. The vegetation communities have been dominated by non-native invasive species and pioneering native speices.

The first section of the report provides the results of the overall morphological assessment of the stream. Primary data products include: a contour (bathymetric) map of the stream, area, volume and depth statistics, and the water level at the time of assessment. These data are useful for evaluating trends and for developing management actions such as plant management where depth and stream volume are needed.

The second section provides the results of the vegetation assessment conducted on the stream. These results can be used to better understand and manage vegetation in the stream. A list is provided with the different plant species found at various sites along the stream. Potentially invasive, exotic (non-native) species are identified in a plant list and the percent of exotics is presented in a summary table. Watershed values provide a means of reference.

The third section provides the results of the water quality sampling of the stream. Both field data and laboratory data are presented. The water quality index (WQI)ⁱ is used to develop a general stream health statement, which is calculated for both the water column with vegetation and the water column if vegetation were removed. These data are derived from the water chemistry and vegetative submerged biomass assessments and are useful in understanding the results of certain stream vegetation management practices.

The intent of this assessment is to provide a starting point from which to track changes in the stream, and where previous comprehensive assessment data is available, to track changes in the stream's general health. These data can provide the information needed to determine changes and to monitor trends in physical condition and ecological health of the stream.

Section 1: Stream Morphology

Bathymetric Mapⁱⁱ. Table 1 provides the stream's morphologic parameters in various units. The bottom of the stream was mapped using a Lowrance HDS 5 with Wide Area Augmentation System (WAAS)ⁱⁱⁱ enabled Global Positioning System (GPS) with fathometer (bottom sounder) to determine the boat's position, and bottom depth in a single measurement. The result is an estimate of the stream's area, mean and maximum depths, and volume and the creation of a bottom contour map (Figure 2). Besides pointing out the deeper fishing holes in the stream, the morphologic data derived from this part of the assessment can be valuable to overall management of the stream vegetation as well as providing flood storage data for flood models.

ⁱ The water quality index is used by the Water Atlas to provide the public with an estimate of their stream resource quality. For more information, see end note 1.

ⁱⁱ A bathymetric map is a map that accurately depicts all of the various depths of a water body. An accurate bathymetric map is important for effective herbicide application and can be an important tool when deciding which form of management is most appropriate for a water body. Stream volumes, hydraulic retention time and carrying capacity are important parts of stream management that require the use of a bathymetric map.

ⁱⁱⁱ WAAS is a form of differential GPS (DGPS) where data from 25 ground reference stations located in the United States receive GPS signals from GPS satellites in view and retransmit these data to a master control site and then to geostationary satellites. For more information, see end note 2.

Table 1. Stream Morphologic Data (Area, Depth and Volume)

Parameter	Feet	Meters	Acres	Acre-Ft	Gallons
Surface Area (sq)	1,326,165	123,205	30.45	0	0
Mean Depth	2.18	0.66	0	0	0
Maximum Depth	7.68	2.34	0	0	0
Volume (cubic)	1,656,031	46,894	0	38.02	12,388,055
Gauge (relative)	11.75	3.58	0	0	0




Figure 2. Overview of the 2014 1-Foot Bathymetric Contour Map for Delaney Creek




Figure 3 2014 1-Foot Bathymetric Contour Map for Dealney Creek Figure 1




Figure 4 2014 1-Foot Bathymetric Contour Map for Dealney Creek Figure 2




Figure 5 2014 1-Foot Bathymetric Contour Map for Dealney Creek Figure 3



Figure 6 2014 1-Foot Bathymetric Contour Map for Dealney Creek Figure 4




Figure 7 2014 1-Foot Bathymetric Contour Map for Dealney Creek Figure 5



Figure 8 2014 1-Foot Bathymetric Contour Map for Dealney Creek Figure 6



Figure 9 2014 1-Foot Bathymetric Contour Map for Dealney Creek Figure 7



Figure 10 2014 1-Foot Bathymetric Contour Map for Dealney Creek Figure 8

Section 2: Stream Ecology (Vegetation)

The stream's apparent vegetative cover and shoreline detail are evaluated using the latest stream aerial photograph as shown in and by use of WAAS-enabled GPS. Submerged vegetation is determined from the analysis of bottom returns from the Lowrance HDS 5 combined GPS/fathometer described earlier. As depicted in **Error! Reference source not found.**¹¹ through 13, 43 vegetation regions have been assessed for in ~250 meter regions measured from the center of the stream. The vegetation assessment regions are set up from the downstream extent and work to the upstream extent. The region beginning and ending points are set using GPS and then loaded into a GIS mapping program (ArcGIS) for display. Each region is sampled in the three primary vegetative zones (emergent, submerged and floating)^{iv}. The latest high resolution aerial photos are used to provide shore details (docks, structures, vegetation zones) and to calculate the extent of surface vegetation coverage. The primary indices of submerged vegetation cover and biomass for the stream, percent area coverage (PAC) and percent volume inhabited (PVI), are determined by transiting the stream by boat and employing a fathometer to collect "hard and soft return" data. These data are later analyzed for presence and absence of vegetation and to determine the height of vegetation if present. The PAC is determined from the presence and absence analysis of 100 sites in the stream and the PVI is determined by measuring the difference between hard returns (stream bottom) and soft returns (top of vegetation) for sites (within the 100 analyzed sites) where plants are determined present.

The data collected during the site vegetation sampling include vegetation type, exotic vegetation, predominant plant species and submerged vegetation biomass. The total number of species from all sites is used to approximate the total diversity of aquatic plants and the percent of invasive-exotic plants on the stream (

^{iv} See end note 3.

Table 2). The Watershed value in Table 2 only includes lakes and streams sampled during the lake and stream assessment project begun in May of 2006. These data will change as additional lakes and streams are sampled.

Figure 12 2014 Delaney Creek Vegetation Assessment Region Map Figure 1

Table 3 through Table 5 detail the results from the 2014 aquatic plant assessment for the stream. These data are determined from the 43 sites used for intensive vegetation surveys. The tables are divided into Floating Leaf, Emergent and Submerged plants and contain the plant code, species, common name and presence (indicated by a 1) or absence (indicated by a blank space) of species and the calculated percent occurrence (number sites species is found/number of sites) and type of plant (Native, Non-Native, Invasive, Pest). In the "Type" category, the codes N and E0 denote species native to Florida. The code E1 denotes Category I invasive species, as defined by the [Florida Exotic Pest Plant Council](#) (FLEPPC); these are species "that are altering native plant communities by displacing native species, changing community structures or ecological functions, or hybridizing with natives." The code E2 denotes Category II invasive species, as defined by FLEPPC; these species "have increased in abundance or frequency but have not yet altered Florida plant communities to the extent shown by Category I species." Use of the term invasive indicates the plant is commonly considered invasive in this region of Florida. The term "pest" indicates a plant (native or non-native) that has a greater than 55% occurrence in the stream and is also considered a problem plant for this region of Florida, or is a non-native invasive that is or has the potential to be a problem plant in the stream and has at least 40% occurrence. These two terms are somewhat subjective; however, they are provided to give stream property owners some guidance in the management of plants on their property. Please remember that to remove or control plants in a wetland (stream shoreline) in Hillsborough County the property owner must secure an [Application To Perform Miscellaneous Activities In Wetlands](#) permit from the [Environmental Protection Commission of Hillsborough County](#) and for management of in-stream vegetation outside the wetland fringe (for streams with an area greater than ten acres), the property owner must secure a [Florida Department of Environmental Protection Aquatic Plant Removal Permit](#).

Table 2. Total Diversity, Percent Exotics, and Number of Pest Plant Species

Parameter	Lake	Watershed
Number of Vegetation Assessment Sites	43	158
Total Plant Diversity (# of Taxa)	109	160
% Non-Native Plants	43	27
Total Pest Plant Species	11	13




Figure 11 2014 Delaney Creek Vegetation Assessment Region Map Overview



Figure 12 2014 Delaney Creek Vegetation Assessment Region Map Figure 1




Figure 13 2014 Delaney Creek Vegetation Assessment Region Map Figure 2

Table 3. List of Floating Leaf Zone Aquatic Plants Found

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
LEN	<i>Lemna spp.</i>	Duckweed	53%	N, E0
ECS	<i>Eichhornia crassipes</i>	Water Hyacinth	46%	E1, P
SMA	<i>Salvinia minima</i>	Water Spangles, Water Fern	46%	E1, P
SPI	<i>Spirodela polyrhiza</i>	Giant Duckweed	39%	N, E0
ACA	<i>Azolla filiculoides</i>	Carolina Mosquito Fern; American Waterfern	25%	N, E0
LSA	<i>Limnobium spongia</i>	American Spongeplant, Frog's Bit	18%	N, E0



Figure 14. Water Hyacinth, *Eichhornia crassipes*, was a common non-native invasive species in the freshwater portions of the Delaney Creek study area.

Table 4. List of Emergent Zone Aquatic Plants Found

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
STS	<i>Schinus terebinthifolius</i>	Brazilian Pepper	83%	E1, P
BMA	<i>Urochloa mutica</i>	Para Grass	65%	E1, P
BAA	<i>Bidens alba</i>	White Beggar-ticks, Romerillo	55%	N, E0
LPA	<i>Ludwigia peruviana</i>	Peruvian Primrosewillow	55%	E1, P
QLA	<i>Quercus laurifolia</i>	Laurel Oak; Diamond Oak	55%	N, E0
PCA	<i>Pontederia cordata</i>	Pickrel Weed	53%	N, E0
ULA	<i>Urena lobata</i>	Caesar's-weed	51%	E1, P
HYE	<i>Hydrocotyle umbellata</i>	Manyflower Marshpennywort, Water Pennywort	48%	N, E0
COM	<i>Commelina spp.</i>	Dayflower	48%	N, E0
CEA	<i>Colocasia esculenta</i>	Wild Taro	46%	E1, P
APS	<i>Alternanthera philoxeroides</i>	Alligator Weed	46%	E2, P
LEL	<i>Leucaena leucocephala</i>	White Leadtree	46%	E2, P
LLA	<i>Ludwigia leptocarpa</i>	Anglestem Primrosewillow	46%	N, E0
EUP	<i>Eupatorium capillifolium</i>	Dog Fennel	41%	N, E0
DBA	<i>Dioscorea bulbifera</i>	Air Potato	39%	E1
SLT	<i>Sagittaria latifolia</i>	Wapato, Common Arrowhead, Broadleaf Arrowhead, Duck Potato	37%	N, E0
AGS	<i>Andropogon glomeratus</i>	Bushy Bluestem; Bush Broom Grass	30%	N, E0
PRS	<i>Panicum repens</i>	Torpedo Grass	30%	E1
MAM	<i>Myriophyllum aquaticum</i>	Parrot Feather	27%	E0

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
ADM	<i>Acrostichum danaeifolium</i>	Giant Leather Fern	27%	N, E0
SLA	<i>Sagittaria lancifolia</i>	Duck Potato	27%	N, E0
RZM	<i>Rhizophora mangle</i>	Red Mangrove	25%	N, E0
MSS	<i>Mikania scandens</i>	Climbing Hempvine	25%	N, E0
PMM	<i>Panicum maximum</i>	Guineagrass	25%	E0
LAG	<i>Laguncularia racemosa</i>	White Mangrove	25%	N, E0
HAS	<i>Hymenachne amplexicaulis</i>	Trompetilla	23%	E1
EAA	<i>Eclipta alba</i>	Yerba De Tajo	20%	N, E0
CYO	<i>Cyperus odoratus</i>	Fragrant Flatsedge	20%	N, E0
BHA	<i>Baccharis halimifolia</i>	Groundsel Tree; Sea Myrtle	20%	N, E0
WTA	<i>Sphagneticola trilobata</i>	Creeping Oxeye; Wedelia	20%	E2
RBA	<i>Ruellia simplex</i>	Britton's Wild Petunia	18%	E1
SCA	<i>Salix caroliniana</i>	Carolina Willow	18%	N, E0
RVS	<i>Rumex verticillatus</i>	Swamp Dock	18%	N, E0
AVG	<i>Avicennia germinans</i>	Black Mangrove	18%	N, E0
ACE	<i>Acer rubrum</i>	Southern Red Maple	18%	N, E0
JRO	<i>Juncus roemerianus</i>	Needle Rush, Black Rush	18%	N, E0
PAR	<i>Paspalum repens</i>	Water Paspalum	18%	N, E0
BRP	<i>Broussonetia papyrifera</i>	Paper Mulberry	16%	E2
SCC	<i>Schoenoplectus californicus</i>	Giant Bulrush	16%	N, E0
AAS	<i>Amaranthus australis</i>	Southern Water Hemp	13%	N,

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
				E0
PDF	<i>Polygonum glabrum</i>	Denseflower Knotweed	13%	N, E0
QNA	<i>Quercus nigra</i>	Water Oak	13%	N, E0
PRA	<i>Pluchea baccharis</i>	Rosy Camphorweed	11%	N, E0
ABM	<i>Amaranthus blitum subsp. Emarginatus</i>	Purple Amaranth; Livid Pigweed	11%	E0
ACS	<i>Symphyotrichum carolinianum</i>	Climbing Aster	9%	N, E0
BOC	<i>Boehmeria cylindrica</i>	Bog Hemp, False Nettle	9%	N, E0
CLA	<i>Casuarina equisetifolia</i>	Australian Pine	9%	E1
LOS	<i>Ludwigia octovalvis</i>	Mexican Primrosewillow, Long-stalked Ludwigia	9%	N, E0
LRS	<i>Ludwigia repens</i>	Creeping Primrosewillow, Red Ludwigia	9%	N, E0
VRA	<i>Vitis rotundifolia</i>	Muscadine Grape	9%	N, E0
WAX	<i>Myrica cerifera</i>	Southern Bayberry; Wax Myrtle	9%	N, E0
TYP	<i>Typha spp.</i>	Cattails	9%	N, E0
SPO	<i>Sabal palmetto</i>	Sabal Palm, Cabbage Palm	6%	N, E0
LPG	<i>Ludwigia peploides glabrescens</i>	Floating Primrosewillow	6%	N, E0
LOP	<i>Ludwigia spp.</i>	Water Primroses, Primrosewillow	6%	E0
ICA	<i>Imperata cylindrica</i>	Cogon Grass	6%	E1
POL	<i>Polygonum spp.</i>	Smartweed, Knotweed	6%	N, E0
PHS	<i>Polygonum hydropiperoides</i>	Mild Waterpepper; Swamp Smartweed	6%	N, E0
OCA	<i>Osmunda cinnamomea</i>	Cinnamon Fern	6%	N, E0

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
BLS	<i>Blechnum serrulatum</i>	Swamp fern, Toothed Midsorus Fern	6%	N
DVA	<i>Diodia virginiana</i>	Buttonweed	6%	N, E0
CAA	<i>Centella asiatica</i>	Asian Pennywort, Coinwort, Spadeleaf	6%	N, E0
CAM	<i>Crinum americanum</i>	Swamp lily	6%	N, E0
BID	<i>Bidens spp.</i>	Beggarticks	4%	E0
EUT	<i>Eustachys petraea</i>	Pinewoods Fingergrass	4%	N, E0
EWI	<i>Echinochloa walteri</i>	Coast Cockspur Grass (hairy)	4%	N, E0
CJE	<i>Cladium jamaicense</i>	Jamaica Swamp Saw Grass	4%	N, E0
PLU	<i>Pluchea spp.</i>	Marsh Fleabane, Camphorweed	4%	N, E0
LJM	<i>Lygodium japonicum</i>	Japanese Climbing Fern	4%	E1
MAH	<i>Melia azedarach</i>	Chinaberry tree	4%	E2
SET	<i>Setaria spp.</i>	Bristlegrass, Foxtail	4%	
SAC	<i>Sacciolepis striata</i>	American Cupscale	4%	N, E0
THA	<i>Thelypteris spp.</i>	Shield ferns	4%	N, E0
UAA	<i>Ulmus americana</i>	American Elm; Florida Elm	2%	N, E0
SAM	<i>Sambucus nigra subsp. Canadensis</i>	Elderberry	2%	N, E0
RCS	<i>Ricinus communis</i>	Castorbean	2%	E2
SHA	<i>Sesbania herbacea</i>	Danglepod Sesban	2%	N, E0
SSM	<i>Sapium sebiferum</i>	Chinese Tallow Tree	2%	E1
SMI	<i>Smilax spp.</i>	Catbriar, Greenbriar	2%	N, E0
SPA	<i>Spartina spp.</i>	Cordgrass	2%	N, E0
SPB	<i>Spartina bakeri</i>	Sand Cordgrass	2%	N,

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
				E0
SPM	<i>Syngonium podophyllum</i>	Nephtytis, Arrowhead Vine, American Evergreen	2%	E1
JUM	<i>Juncus marginatus</i>	Shore Rush, Grassleaf Rush	2%	N, E0
IRI	<i>Iris spp.</i>	Flag	2%	E0
ITE	<i>Itea virginica</i>	Virginia Willow; Virginia Sweetspire	2%	N, E0
IVA	<i>Iris virginica</i>	Southern Blue Flag	2%	N, E0
PQA	<i>Parthenocissus quinquefolia</i>	Virginia Creeper, Woodbine	2%	N, E0
NEA	<i>Nephrolepis exaltata</i>	Sword Fern, Wild Boston Fern	2%	N, E0
PFO	<i>Paederia foetida</i>	Skunkvine, Stinkvine	2%	E1
PHN	<i>Panicum hemitomon</i>	Maidencane	2%	N, E0
CER	<i>Ceratopteris thalictroides</i>	Water Sprite	2%	E0
CLG	<i>Cyperus ligularis</i>	Swamp Flatsedge	2%	N, E0
CMA	<i>Cicuta maculata var. mexicana</i>	Spotted Water Hemlock	2%	N, E0
CPT	<i>Cyperus polystachyos</i>	Flat Sedge	2%	N, E0
CSS	<i>Cyperus surinamensis</i>	Tropical Flatsedge	2%	N, E0
CYP	<i>Cyperus spp.</i>	Sedge	2%	E0
CCA	<i>Cinnamomum camphora</i>	Camphor-tree	2%	E1



Figure 15. The emergent vegetation zone of the freshwater portion of Delaney Creek was dominated by Paragrass, *Urochloa mutica*.



Figure 16 The tidal portions of the Delaney Creek study area was dominated by mangroves, *laguncaria racemosa* and *rhizophora mangle*, and Brazilian Pepper, *schinus terebinthifolous*.

Table 5. List of Submerged Zone Aquatic Plants Found.

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
HPA	<i>Hygrophila polysperma</i>	East Indian Hygrophila, Indian Swampweed	51%	E1, P
HVA	<i>Hydrilla verticillata</i>	Hydrilla, waterthyme	51%	E1, P
LSE	<i>Limnophila sessiliflora</i>	Asian Marshweed	18%	E2
CDM	<i>Ceratophyllum demersum</i>	Hornwort, Coontail	13%	N, E0
BMI	<i>Bacopa monnieri</i>	Common Bacopa	11%	N, E0



Figure 17. Hydrilla, *hydrilla verticillata*, was a common non-native invasive species dominating the submerged vegetation zone of the freshwater portions of the Delaney Creek study area.

Table 6. List of All Plants and Sample Sites

Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
Brazilian Pepper	1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,27,34,35,36,37,38,39,40,41,42,43	83	Emergent
Para Grass	16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43	65	Emergent
Laurel Oak; Diamond Oak	14,15,17,18,19,20,21,22,26,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43	55	Emergent
Peruvian Primrosewillow	18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,35,37,38,39,40,41,42,43	55	Emergent
White Beggar- ticks, Romerillo	19,20,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43	55	Terrestrial
Duckweed	16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,38,39,40,41,42,43	53	Floating
Pickrel Weed	18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,37,38,39,40,41,42	53	Emergent
Caesar's-weed	18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,38,42,43	51	Emergent
East Indian Hygrophila, Indian Swampweed	22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43	51	Submersed
Hydrilla, waterhyme	22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43	51	Submersed
Dayflower	22,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43	48	Emergent
Manyflower Marshpennywort, Water Pennywort	16,17,18,19,20,21,22,23,24,25,26,27,35,36,37,38,39,40,41,42,43	48	Emergent
Alligator Weed	14,15,16,17,18,19,20,21,22,23,24,25,26,27,35,39,40,41,42,43	46	Emergent
Anglestem Primrosewillow	22,23,24,25,26,27,28,29,30,31,32,33,34,37,38,39,40,41,42,43	46	Emergent
Water Hyacinth	16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35	46	Floating
Water Spangles, Water Fern	14,15,16,17,18,19,20,21,22,23,24,25,26,37,38,39,40,41,42,43	46	Floating
White Leadtree	8,9,12,13,14,15,16,18,19,20,21,22,23,27,37,38,39,41,42,43	46	Terrestrial
Wild Taro	18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37	46	Emergent
Dog Fennel	19,20,22,23,24,25,26,27,32,33,34,37,38,39,40,41,42,43	41	Emergent
Air Potato	14,15,16,17,18,19,20,21,22,23,29,30,31,32,33,35,43	39	Emergent
Giant Duckweed	22,23,24,25,26,27,28,29,30,31,32,38,39,40,41,42,43	39	Floating
Wapato,	18,19,20,22,25,26,27,28,29,30,31,32,33,39,41,42	37	Emergent

Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
Common Arrowhead, Broadleaf Arrowhead, Duck Potato			
Bushy Bluestem; Bush Broom Grass	22,23,24,25,26,27,28,29,30,31,32,33,34	30	Emergent
Torpedo Grass	18,19,22,23,24,26,30,38,39,40,41,42,43	30	Emergent
Duck Potato	18,19,22,23,24,25,27,28,29,30,31,32	27	Emergent
Giant Leather Fern	6,12,13,14,15,16,17,18,19,20,21,22	27	Emergent
Parrot Feather	22,23,24,25,26,27,28,29,30,31,32,33	27	Emergent
Carolina Mosquito Fern; American Waterfern	22,23,24,25,26,27,28,29,30,31,32	25	Floating
Climbing Hempvine	22,23,24,25,26,27,28,29,30,31,32	25	Emergent
Guineagrass	16,17,18,19,24,25,39,40,41,42,43	25	Terrestrial
Red Mangrove	1,2,3,4,5,6,7,8,9,10,11	25	Terrestrial
White Mangrove	1,2,3,4,5,6,7,8,9,10,11	25	Terrestrial
Trompetilla	24,25,26,27,28,29,30,31,32,35	23	Emergent
Creeping Oxeye; Wedelia	20,23,24,30,34,40,41,42,43	20	Emergent
Fragrant Flatsedge	23,24,26,27,39,40,41,42,43	20	Emergent
Groundsel Tree; Sea Myrtle	22,25,28,29,30,31,32,33,34	20	Emergent
Yerba De Tajo	18,19,23,24,30,31,32,42,43	20	Emergent
American Spongeplant, Frog's Bit	14,15,16,17,18,19,20,21	18	Floating
Asian	14,15,16,17,18,19,20,21	18	Submersed

Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
Marshweed			
Black Mangrove	1,2,3,4,5,6,7,8	18	Terrestrial
Britton's Wild Petunia	18,19,20,21,22,23,24,25	18	Terrestrial
Carolina Willow	18,19,20,22,23,24,30,42	18	Emergent
Needle Rush, Black Rush	9,10,11,12,13,17,18,25	18	Emergent
Southern Red Maple	22,23,33,34,37,39,40,42	18	Emergent
Swamp Dock	14,17,18,19,20,21,22,35	18	Emergent
Water Paspalum	25,28,29,30,31,32,33,37	18	Emergent
Giant Bulrush	18,19,20,21,22,23,24	16	Emergent
Paper Mulberry	30,38,39,40,41,42,43	16	Emergent
Denseflower Knotweed	22,23,24,25,26,27	13	Emergent
Hornwort, Coontail	22,23,24,25,26,27	13	Submersed
Southern Water Hemp	16,17,18,19,20,42	13	Emergent
Water Oak	16,22,23,33,38,39	13	Emergent
Common Bacopa	13,22,25,40,43	11	Submersed
Purple Amaranth; Livid Pigweed	16,17,18,19,20	11	Emergent
Rosy Camphorweed	17,19,22,23,43	11	Emergent
Australian Pine	10,11,12,13	9	Emergent
Bog Hemp, False Nettle	14,18,19,41	9	Emergent
Cattails	17,19,20,28	9	Emergent
Climbing Aster	20,23,24,28	9	Emergent
Creeping Primrosewillow, Red Ludwigia	22,25,26,39	9	Emergent

Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
Mexican Primrosewillow, Long-stalked Ludwigia	19,24,25,26	9	Emergent
Muscadine Grape	37,38,42,43	9	Emergent
Southern Bayberry; Wax Myrtle	27,33,38,43	9	Emergent
Asian Pennywort, Coinwort, Spadeleaf	22,23,35	6	Emergent
Buttonweed	23,24,25	6	Emergent
Cinnamon Fern	21,22,35	6	Emergent
Cogon Grass	23,27,34	6	Terrestrial
Floating Primrosewillow	18,19,20	6	Emergent
Mild Waterpepper; Swamp Smartweed	21,40,42	6	Emergent
Sabal Palm, Cabbage Palm	22,23,25	6	Terrestrial
Smartweed, Knotweed	18,19,20	6	Emergent
Swamp fern, Toothed Midsorus Fern	21,22,23	6	Emergent
Swamp lily	17,18,22	6	Emergent
Water Primroses, Primrosewillow	18,19,20	6	Emergent
American Cupscale	33,34	4	Emergent

Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
Beggarticks	19,20	4	Emergent
Bristlegrass, Foxtail	8,9	4	Emergent
Chinaberry tree	15,18	4	Emergent
Coast Cockspur Grass (hairy)	19,21	4	Emergent
Jamaica Swamp Saw Grass	17,18	4	Emergent
Japanese Climbing Fern	30,35	4	Terrestrial
Marsh Fleabane, Camphorweed	17,19	4	Emergent
Pinewoods Fingergrass	42,43	4	Terrestrial
Shield ferns	35,36	4	Emergent
American Elm; Florida Elm	23	2	Emergent
Aster spp., Elliot's Aster	20	2	Unknown
Camphor-tree	21	2	Emergent
Castorbean	37	2	Terrestrial
Catbriar, Greenbriar	21	2	Emergent
Chinese Tallow Tree	42	2	Emergent
Cordgrass	18	2	Emergent
Danglepod Sesban	26	2	Emergent
Elderberry	23	2	Emergent
Flag	18	2	Emergent
Flat Sedge	24	2	Emergent
Maidencane	28	2	Emergent
Nephthytis, Arrowhead Vine,	37	2	Terrestrial

Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
American Evergreen			
Sand Cordgrass	18	2	Terrestrial
Sedge	19	2	Emergent
Shore Rush, Grassleaf Rush	19	2	Emergent
Skunkvine, Stinkvine	35	2	Terrestrial
Southern Blue Flag	22	2	Emergent
Spotted Water Hemlock	19	2	Emergent
Swamp Flatsedge	42	2	Terrestrial
Sword Fern, Wild Boston Fern	22	2	Terrestrial
Tropical Flatsedge	22	2	Emergent
Virginia Creeper, Woodbine	16	2	Emergent
Virginia Willow; Virginia Sweetspire	21	2	Emergent
Water Sprite	18	2	Emergent

Discussion of Vegetation Assessment Results

The highest diversity of vegetation in the Delaney Creek study area was found in the freshwater portions where several species of non-native invasive species were found growing alongside pioneer native species in disturbed areas. The lowest diversity of vegetation was found in the tidal regions near the mouth where the influence of salinity and daily fluctuating water levels left only those species able to cope with such stresses.

Section 3: Long-term Ambient Water Chemistry

A critical element in any stream assessment is the long-term water chemistry data set. These data are obtained from several data sources that are available to the Water Atlas and are managed in the Water Atlas Data Download and graphically presented on the water quality page for streams in Hillsborough County. The Delaney Creek Water Quality Page can be viewed at <http://www.hillsborough.wateratlas.usf.edu/river/waterquality.asp?wbodyid=27&wbodyatlas=river>.

A primary source of stream water chemistry in Hillsborough County is the Routine Monitoring Sampling by the Hillsborough County Environmental Protection Commission. Other source data are used as available; however these data can only indicate conditions at time of sampling.

These data are displayed and analyzed on the Water Atlas as shown in Figure 18, Figure 19, and Figure 20 for Delaney Creek. The figures are graphs of: (1) the overall water quality index (WQI), which is a method commonly used to characterize the productivity of a stream, and may be thought of as a stream's ability to support plant growth and a healthy food source for aquatic life; (2) the chlorophyll *a* concentration, which indicates the stream's algal concentration, and (3) the stream's Secchi Disk depth which is a measure of water visibility and depth of light penetration. These data are used to evaluate a stream's ecological health and to provide a method of ranking streams and are indicators used by the US Environmental Protection Agency (USEPA) and the Florida Department of Environmental Protection (FDEP) to determine a stream's level of impairment. The chlorophyll *a* and Secchi Disk depth graphs include benchmarks which indicate the median values for the various parameters for a large number of Streams in Florida expressed as percentiles.

Based on best available data, Delaney Creek has a color value determined as a platinum cobalt unit (pcu) value of 44.0 at 36th Avenue, 10.2 at the mouth and is considered a Dark stream (has a mean color in pcu greater than 40 in the freshwater portions). The FDEP and USEPA may classify a stream as impaired if the stream is a dark stream and has a WQI greater than 60, or is a clear stream (has a mean color in pcu less than or equal to 40) and has a WQI greater than 40. Delaney Creek has a WQI of 21 and does not meet the FDEP Impaired Waters Rule (IWR) criteria for impaired streams. See also Table .




Figure 18. Recent Water Quality Index (WQI) graph for Delaney Creek^v

^v Graph source: Hillsborough County Water Atlas. For an explanation of the Good, Fair and Poor benchmarks, please see the notes at the end of this report. For the latest data go to: http://www.hillsborough.wateratlas.usf.edu/graphs20/graph_it.aspx?wbodyid=27&data=WQI&data_type=WQ&waterbodyatlas=river&ny=10&bench=1




Figure 19. Recent Chlorophyll a graph for Delaney Creek^{vi}

^{vi} Graph Source: Hillsborough County Water Atlas. For the latest data go to http://www.hillsborough.wateratlas.usf.edu/graphs20/graph_it.aspx?wbodyid=27&data=Chla_ugl&datatype=WQ&waterbodyatlas=river&ny=10&bench=1




Figure 20. Recent Secchi Disk graph for Delaney Creek^{vii}

^{vii} Graph Source: Hillsborough County Water Atlas. For the latest data go to http://www.hillsborough.wateratlas.usf.edu/graphs20/graph_it.aspx?wbodyid=27&data=secchi_ft&datatype=WQ&waterbodyatlas=stream&ny=10&bench=1

Stream Numeric Nutrient Criteria. November 30, 2012 the USEPA accepted the majority of the FDEP proposed NNCs which included an NNC for streams. The NNC for freshwater streams is provided in the Stream Assessment Notes at the end of this report, and for the Tampa Bay area (considered West Central) total phosphorous must be less than or equal to 0.49 mg/L and total nitrogen must be less than or equal to 1.65 mg/L to meet the criteria (Table 7) and chlorophyll a must be at or below 20 µg/L not be considered impaired.

Table 7 Stream Numeric Nutrient Criteria

<u>Nutrient Watershed Region</u>	<u>Total Phosphorus Nutrient Threshold¹</u>	<u>Total Nitrogen Nutrient Threshold¹</u>
<u>Panhandle West</u>	<u>0.06 mg/L</u>	<u>0.67 mg/L</u>
<u>Panhandle East</u>	<u>0.18 mg/L</u>	<u>1.03 mg/L</u>
<u>North Central</u>	<u>0.30 mg/L</u>	<u>1.87 mg/L</u>
<u>Peninsular</u>	<u>0.12 mg/L</u>	<u>1.54 mg/L</u>
<u>West Central</u>	<u>0.49 mg/L</u>	<u>1.65 mg/L</u>
<u>South Florida</u>	<u>No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.</u>	<u>No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.</u>

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year period.

Delaney Creek, a freshwater creek that flows into Hillsborough Bay, has four long-term data stations in the study area (Hillsborough County EPC Routine Monitoring 133 and 138, Quarterly 606 and 607) whose three-year geometric mean for Total Nitrogen, Total Phosphorus and Chlorophyll are as shown below in Table 8. According to the Numeric Nutrient Criteria, Delaney Creek does not exceed the numeric nutrient criteria for Phosphorous, Nitrogen or Chlorophyll concentrations.

Table 8 Flint Creek NNC data summary

Delaney Creek Study Area	Total Phosphorous mg/l	Total Nitrogen mg/l	Chlorophyll-a Corrected µg/l
Period of Record Geomean	0.461	1.794	4.19
2012 Geomean	0.220	1.242	5.14
2013 Geomean	0.206	1.183	5.24
2014 Geomean	0.228	0.967	6.39

As part of the stream assessment the physical water quality and chemical water chemistry of a stream are measured. These data only indicate a snapshot of the stream's water quality; however they are useful when compared to the trend data available from Hillsborough County Environmental Protection Commission or other sources. Table 9 contains the summary water quality data and index values and adjusted values calculated from these data. The total phosphorus (TP), total nitrogen (TN) and chlorophyll a water chemistry sample data are the results of chemical analysis of samples taken during the assessment and analyzed by the Hillsborough County Environmental Protection Commission laboratory.

The growth of plants (planktonic algae, macrophytic algae and rooted plants) is directly dependent on the available nutrients within the water column of a stream and to some extent the nutrients which are held in the sediment and the vegetation biomass of a stream. Additionally, algae and other plant growth are [limited](#) by the nutrient in lowest concentration relative to that needed by a plant. Plant biomass contains less phosphorus by weight than nitrogen so phosphorus is many times the limiting nutrient. When both nutrients are present at a concentration in the stream so that either or both may restrict plant growth, the limiting factor is called "balanced". The ratio of total nitrogen to total phosphorus, the "N to P" ratio (N/P), is used to determine the limiting factor. If N/P is greater than or equal to 30, the stream is considered phosphorus limited, when this ratio is less than or equal to 10, the stream is considered nitrogen limited and if between 10 and 30 it is considered balanced.

Table 9. Water Quality Parameters (Laboratory) for Delaney Creek

Parameter	Mouth	36 th Ave S
Total Phosphorus (ug/L)	241	202
Total Nitrogen (ug/L)	420	750
Chlorophyll a Corrected(ug/L)	10.3	16.2
TN/TP	1.74	3.71
Limiting Nutrient	Nitrogen	Nitrogen
Color (PCU)	10.2	44.0
Secchi disk depth (ft)	4.9	2.5

The color of a stream is also important to the growth of algae. Dark, tannic streams tend to suppress algal growth and can tolerate a higher amount of nutrient in their water column; while clear streams tend to support higher algal growth with the same amount of nutrients. The color of a stream, which is measured in a unit called the "cobalt platinum unit (PCU)" because of the standard used to determine color, is important because it is used by the State of Florida to determine stream impairment as explained earlier. Rivers, streams or other "flow through" systems tend to support lower algal growth for the same amount of nutrient concentration. All these factors are important to the understanding of your stream's overall condition. Table 10 includes many of the factors that are typically used to determine the actual state of plant growth in your stream. These data should be understood and reviewed when establishing a management plan for a stream; however, as stated above other factors must be considered when developing such a plan. Please contact the [Water Atlas Program](#) if you have questions about this part or any other part of this report.

Table 10 provides data derived from the vegetation assessment which is used to determine an adjusted WQI. This is accomplished by calculating the amount of phosphorus and nitrogen that could be released by existing submerged vegetation (Adjusted Nutrient) if this vegetation were treated with an herbicide or managed by the addition of Triploid Grass Carp (*Ctenopharyngodon idella*). The table also shows the result of a model that calculates the potential algae, as chlorophyll a (Adjusted Chlorophyll), which could develop due to the additional nutrients held within the plant biomass. While it would not be expected that all the vegetation would be turned into available phosphorus by these management methods, the data is useful when planning various management activities. Approximately 9.0 % of the stream has submerged vegetation present (PAC) and this vegetation represents about 2.5 % of the available stream volume (PVI). Please see additional parameters for adjusted values where appropriate in Table 10. The

vegetation holds enough nutrients to add about 207 $\mu\text{g/L}$ of phosphorus and 7,040 $\mu\text{g/L}$ of nitrogen to the water column and increase the algal growth potential within the stream.

Delaney Creek is nitrogen-limited; i.e., an increase in nitrogen could change the WQI and increase the potential for algal growth.

Table 10. Field parameters and calculations used to determine nutrients held in Submerged Aquatic Vegetation (SAV) biomass.

Parameter	Value
% Area Covered (PAC)	2.10 %
PVI	0.51 %
Total Phosphorus - Adjusted (ug/L)	207
Total Phosphorus - Combined (ug/L)	410
Total Nitrogen - Adjusted (ug/L)	7,040
Total Nitrogen - Combined (ug/L)	7,790
Chlorophyll - Adjusted from Total Nutrients (ug/L)	18.28
Chlorophyll - Combined (ug/L)	54.58

Table 1111 contains the field data taken in the upstream and downstream extents of the stream using a multi-probe (we use a Eureka Manta) which has the ability to directly measure the temperature, pH, dissolved oxygen (DO), percent DO (calculated from DO, temperature and conductivity). These data are listed for three levels in the stream.

Table 11. Water Chemistry Data Based on Manta Water Chemistry Probe for Delaney Creek

Sample Location	Sample Depth (m)	Time	Temp (deg C)	Conductivity (mS/cm3)	Dissolved Oxygen (%)	Dissolved Oxygen (mg/L)	pH
Mean Value	1.48	10/25/2013 12:00:00 AM	23.99	0.426	94.16	8.31	8.01
Surface - 78th Street	0.91	10/25/2013 12:00:00 AM	23.83	0.383	86.60	7.68	8.08
Surface - 86th Street	1.27	10/25/2013 12:00:00 AM	23.68	0.360	118.60	10.54	8.15
Surface - NS canal Robindale Road	0.93	10/25/2013 12:00:00 AM	23.68	0.483	66.90	5.94	7.60
Surface - 86th Street	0.28	7/22/2014 12:00:00 AM	28.76	0.168	80.30	6.23	7.24
Bottom - 86th Street	0.88	7/22/2014 12:00:00 AM	28.76	0.168	76.50	5.94	7.15
Surface - Mouth	0.47	7/22/2014 12:00:00 AM	28.72	0.742	69.58	5.39	7.16

To better understand many of the terms used in this report, we recommend that the reader visit the [Hillsborough County & City of Tampa Water Atlas](#) and explore the “Learn More” areas which are found on the resource pages. Additional information can also be found using the [Digital Library](#) on the Water Atlas website.

Section 4: Conclusion

Delaney Creek is a small area (30.4-acre) stream that would be considered in the healthy category of streams based on water chemistry. It has a plant diversity of 109 species relative to the total watershed plant diversity of 160 species with about 9.00 % percent of the open water areas containing submerged aquatic vegetation. Vegetation helps to maintain the nutrient balance in the stream as well as provide good fish habitat. The stream has few open water areas to support various types of recreation and has a poor diversity of plant species. The primary pest plants in the stream include *Schinus terebinthifolius*, *Urochloa mutica*, *Ludwigia peruviana*, *Urena lobata*, *Hygrophila polysperma* and *Hydrilla verticillata*.

This assessment was accomplished to assist stream property owners to better understand and manage their streams. Hillsborough County supports this effort as part of their [Stream Waterwatch Program \(SWW\)](#) and has developed guidelines for stream property owner groups to join the SWW and receive specific assistance from the County in the management of their stream. For additional information and recent updates please visit the [Hillsborough County & City of Tampa Water Atlas](#) website.

Stream Assessment Notes

1. The Water Quality Index (WQI)¹ is similar to the Trophic State Index (TSI) in that both are used for the statewide assessment of surface waters: the 305(b) Report. WQI is used for streams, black waters (natural tea and coffee-colored waters), and springs, while TSI is used for lakes and estuaries. The WQI is calculated by averaging the values of most or all of the parameters within five water quality parameter categories: 1) water clarity (measured as turbidity and-or Secchi disk depth), 2) dissolved oxygen, 3) oxygen demanding substances (measured as biochemical oxygen, chemical oxygen demand and-or total organic carbon), 4) nutrients (measured as total nitrogen, nitrite plus nitrate, and-or total phosphorus), and 5) bacteria (total coliform and-or fecal coliform).

Water Atlas presents WQIs over the last four seasons (three month intervals). The WQI "value" for a waterbody is determined by averaging the values (data) of the aforementioned parameters for each "season" (Jan-Mar, Apr-Jun, Jul-Sep, Oct-Dec). These seasonal averages are then averaged to provide an overall "rating" or WQI. The term "confidence" expresses the degree of completeness of the index; in other words, "confidence" states how many parameter categories were used to calculate the Overall Water Quality Index.

Ranges of WQI values have been established to provide a general ranking of the waterbody (Figure 1.) WQI values may also include the 'Confidence' (Figure 2) , which provides you with some relative idea as to how much information was used to calculate the WQI for that waterbody.

WQI	Rating
0-45	Good
45-60	Fair
>60	Poor


Figure 1. Water Quality Index (WQI) ranges and their designations.

WQI	Rating	Confidence	Season
30	Good	5/5	Winter (2000)
40	Good	3/5	Fall (2000)
30	Good	2/5	Summer (2000)
50	Fair	3/5	Summer (2000)

Figure 2. WQI rankings are provided with examples of Confidence values.

¹The acronym WQI also stands for "Water Quality Inspection" in much of the DEP literature.

2. **Wide Area Augmentation System (WAAS)** is a form of differential GPS (DGPS) where data from 25 ground reference stations located in the United States receive GPS signals from GPS satellites in view and retransmit these data to a master control site and then to geostationary satellites. The geostationary satellites broadcast the information to all WAAS-capable GPS receivers. The receiver decodes the signal to provide real time correction of raw GPS satellite signals also received by the unit. WAAS-enabled GPS is not as accurate as standard DGPS which employs close by ground stations for correction, however; it was shown to be a good substitute when used for this type of mapping application. Data comparisons were conducted with both types of DGPS employed simultaneously and the positional difference was determined to be well within the tolerance established for the project.
3. The three primary aquatic vegetation zones are shown below:



4. A stream is **impaired** if: "A stream or stream segment shall be included on the planning list for nutrients if the following imbalances are observed:
 - a. Algal mats are present in sufficient quantities to pose a nuisance or hinder reproduction of a threatened or endangered species, or
 - b. Annual mean chlorophyll a concentrations are greater than 20 $\mu\text{g/l}$ or if data indicate annual mean chlorophyll a values have increased by more than 50% over historical values for at least two consecutive years.

Specific Authority 403.061, 403.067 FS. Law Implemented 403.062, 403.067 FS. History – New 6-10-02, Repromulgated 1/2/07."

Please see page 12 of the [Impaired Waters Rule](#). Updated activity regarding impaired waters may be tracked at: <http://www.dep.state.fl.us/water/tmdl/>

5. An **adjusted chlorophyll a value** ($\mu\text{g/L}$) was calculated by modifying the methods of Canfield et al (1983). The total wet weight of plants in the stream (kg) was calculated by multiplying stream surface area (m^2) by PAC (percent area coverage of macrophytes) and multiplying the product by the biomass of submersed plants (kg wet weight m^2) and then by 0.25, the conversion for the 1/4 meter sample cube. The dry weight (kg) of plant material was calculated by multiplying the wet weight of plant material (kg) by 0.08, a factor that represents the average percent dry weight of submersed plants (Canfield and Hoyer, 1992) and then converting to grams. The potential phosphorus concentration (mg/m^3) was calculated by multiplying dry weight (g) by 1.41 mg TP g^{-1} dry weight, a number that represents the mean phosphorus (mg) content of dried plant material measured in 750 samples from 60 Florida lakes (University of Florida, unpublished data), and then dividing by lake volume (m^3) and then converting to $\mu\text{g/L}$ (1000/1000). From the potential phosphorus concentration, a predicted chlorophyll a concentration was determined from the total phosphorus and chlorophyll a relationship reported by Brown (1997) for 209 Florida lakes. Adjusted chlorophyll a concentrations were then calculated by adding each lake's measured chlorophyll a concentration to the predicted chlorophyll a concentration.