

The lake assessments are created in partnership with Hillsborough County and the Florida Center for Community Design and Research

#### LAKE ASSESSMENT DOCUMENT

Lake Helen 8/25/99 Watershed: Rocky/ Brushy Creek

Lake assessments are being conducted to contribute physical and ecological data to the Atlas as a collaborative effort between project partners. The goal is to rapidly assess many of the lakes in the county and thus provide stakeholders a better understanding of the character of the lake, its shore, and the aquatic plants present there. These data are intended to assist in the future management of the lake and its watershed.

The first section of the report provides the results of the bottom mapping effort: a contour (bathymetric) map of the lake, area, volume and depth statistics, and the water level at the time of assessment (if available).

The second section provides the results of the ecological (vegetation) assessment conducted on the lake. These results can be used to better manage vegetation in your lake. A list is provided with the different plant species found at various sites around the lake. Potentially invasive, exotic (non-native) species are identified in a plant list and the percent of exotics is presented in a summary table. The results of this study are compared with other lakes in the watershed.

The intent of the assessment is to provide a starting point from which to track changes in your lake. These data can provide the information needed to determine changes and to monitor trends in physical condition and ecological health of the lake.

### **I. Physical Data** – Area, Depth, Volume, & Bottom Contours

The bottom of the lake was mapped using a sophisticated Global Positioning System (GPS) to determine the boat's position, and a depth-finder to provide depth associated with that measured position. The result is an estimate of your lake's area, mean and maximum depths, and volume (Table 1) and the creation of a bottom contour map.

Table 1. Physical Characteristics of Your Lake.

| Surface Area (acres): | 16         |
|-----------------------|------------|
| Mean Depth (feet):    | 8.9        |
| Maximum Depth (feet): | 25.8       |
| Volume (gallons):     | 47,025,186 |

## Ø SY $\varphi$ 24 S. 46

# Lake Helen

Section - Township - Range 19-27-18



Contour Lines Expressed in 2-Foot Intervals

Lake Perimeter Ground Level


### **EXPLANATION:**

Survey Date August 25, 1999.

surveyed. Contours are expressed above sea level when the lake was in absolute depth below this level. Lake water level was 50 ft.

## DATA SOURCES:

Digital orthophotos by United States Geological Survey. All contours generated by Florida Center for based on survey data provided by the Hillsborough County Lake Management Program. Community Design and Research







The lake assessments are created in partnership with Hillsborough County and the Florida Center for Community Design and Research

#### LAKE ASSESSMENT DOCUMENT

Lake Helen 8/25/99 Watershed: Rocky/ Brushy Creek

### II. Ecological Data

**Aquatic Plant Survey** 

Approximately equispaced sites are haphazardly mapped around the lake and the aquatic plants at each site are surveyed. The total number of species from all sites is used to approximate the total diversity of aquatic plants and the percent of invasive-exotic plants on the lake and in the watershed (Table 2). Many of these plants are considered ecologically harmful, as they tend to out-compete native species. Such "nuisance" plants can also make boating and other recreational activities difficult or impossible. The common and scientific names of plant species found on your lake are listed in Table 3.

Table 2. Comparison of species diversity between your lake and other assessed lakes located within your watershed.

|                        | Lake Helen | Rocky/ Brushy Creek |  |
|------------------------|------------|---------------------|--|
|                        |            | (Average)           |  |
| Number of Taxa:        | 44         | 35                  |  |
| Percent Exotic Plants: | 18%        | 18%                 |  |

Table 3. Botanical and common names of the most commonly found plants on your lake. Percent frequency (of occurence), habit (location where found), status (native or exotic), and EPPC status are provided.

| Common Name                            | Plant Species                     | Frequency | Habit     | Status  | EPPC |
|--|-----------------------------------|-----------|-----------|---------|------|
| Torpedo Grass                          | Panicum repens                    | 100%      | Emergent  | Exotic  | 1    |
| Cypress                                | Taxodium spp.                     | 100%      | Emergent  | Native  | NL   |
| Baldwin's Spikerush, Roadgrass         | Eleocharis baldwinii              | 90%       | Submersed | Native  | NL   |
| Maidencane                             | Panicum hemitomon                 | 90%       | Emergent  | Native  | NL   |
| Pickerel Weed                          | Pontederia cordata                | 90%       | Emergent  | Native  | NL   |
| Creeping Oxeye                         | Sphagneticola (Wedelia) trilobata | 90%       | Emergent  | Exotic  | II   |
| Water Primroses, Primrosewillow        | Ludwigia spp.                     | 70%       | Emergent  | Unknown | NL   |
| Swamp Fern                             | Blechnum serrulatum               | 60%       | Emergent  | Native  | NL   |
| Common Buttonbush                      | Cephalanthus occidentalis         | 60%       | Emergent  | Native  | NL   |
| Southern Red Maple                     | Acer rubrum var. trilobum         | 50%       | Emergent  | Native  | NL   |
| Rush Fuirena                           | Fuirena spp.                      | 50%       | Emergent  | Native  | NL   |
| Manyflower Marshpennywort, Water Penny | Hydrocotyl umbellata              | 50%       | Emergent  | Native  | NL   |
| Loblolly Bay                           | Gordonia lasianthus               | 40%       | Emergent  | Native  | NL   |
| Climbing Hempvine                      | Mikania scandens                  | 40%       | Emergent  | Native  | NL   |
| Fourpetal St. John's-Wort              | Hypericum tetrapetalum            | 30%       | Emergent  | Native  | NL   |
| Punk Tree, Melaleuca                   | Melaleuca quinquenervia           | 30%       | Emergent  | Exotic  | 1    |
|  |                                   |           |           |         |      |

### Lake Helen 8/25/99 Watershed: Rocky/ Brushy Creek

| Laurel Oak; Diamond Oak                    | Quercus laurifolia       | 30% | Emergent  | Native  | NL     |
|--|--------------------------|-----|-----------|---------|--------|
| Algal Mats, Floating                       | Algal spp.               | 20% | Floating  | Unknown | Unknow |
| Pigweed, Water Hemp                        | Amaranthus australis     | 20% | Emergent  | Native  | NL     |
| Common Bacopa, Herb-Of-Grace               | Bacopa monnieri          | 20% | Submersed | Native  | NL     |
| Asian Pennywort, Coinwort, Spadeleaf       | Centella asiatica        | 20% | Emergent  | Native  | NL     |
| Camphor-tree                               | Cinnamomum camphora      | 20% | Emergent  | Native  | 1      |
| Swamp Rosemallow, Swamp Hibiscus           | Hibiscus grandiflorus    | 20% | Emergent  | Native  | NL     |
| Sweetbay Magnolia                          | Magnolia virginiana      | 20% | Emergent  | Native  | NL     |
| Wax Myrtle                                 | Myrica cerifera          | 20% | Emergent  | Native  | NL     |
| Stonewort                                  | Nitella spp.             | 20% | Submersed | Native  | NL     |
| Algae                                      | Periphyton spp.          | 20% | Submersed | Native  | NL     |
| Willow                                     | Salix spp.               | 20% | Emergent  | Native  | NL     |
| Filamentous Algae, Algal Mats              | Algae, Filamentous       | 10% | Floating  | Native  | NL     |
| Wild Taro, Dasheen, Coco Yam               | Colocasia esculenta      | 10% | Emergent  | Exotic  | 1      |
| Dayflower                                  | Commelina diffusa        | 10% | Emergent  | Exotic  | NL     |
| Fragrant Flatsedge                         | Cyperus odoratus         | 10% | Emergent  | Native  | NL     |
| Sedge                                      | Cyperus spp.             | 10% | Emergent  | Unknown | NL     |
| Buttonweed                                 | Diodia virginiana        | 10% | Emergent  | Native  | NL     |
| False Daisy, Yerba De Tajo                 | Eclipta alba (prostrata) | 10% | Emergent  | Native  | NL     |
| Dahoon Holly                               | Ilex cassine             | 10% | Emergent  | Native  | NL     |
| Spatterdock, Yellow Pondlily               | Nuphar lutea var. advena | 10% | Floating  | Native  | NL     |
| Frog-fruit, Carpetweed, Turkey Tangle Fogf | Phyla nodiflora          | 10% | Emergent  | Native  | NL     |
| Pine Tree                                  | Pinus spp.               | 10% | Emergent  | Native  | NL     |
| Marsh Fleabane,Camphorweed                 | Pluchea spp.             | 10% | Emergent  | Native  | NL     |
| Popcorn Tree, Chinese Tallow Tree          | Sapium sebiferum         | 10% | Emergent  | Exotic  | 1      |
| Brazilian Pepper                           | Schinus terebinthifolius | 10% | Emergent  | Exotic  | 1      |
| Cattails                                   | Typha spp.               | 10% | Emergent  | Native  | NL     |
| Elephant-Ear                               | Xanthosoma sagittifolium | 10% | Emergent  | Exotic  | NL     |
|  |                          |     |           |         |        |

### **Standing Crop**

In addition to an overall survey of the types of plants on a lake, an estimate of the standing crop (biomass) of the lake has been obtained for many lakes. This was done by calculating the average weight of the vegetation within a quarter-meter square quadrat tossed haphazardly into three zones (see Figure) at each sampling site around the lake: (1) the emergent zone, (2) the floating zone and (3) the submersed zone. The average weight of the plants (Table 4) from all sampling sites and the dominant type of vegetation (Table 5) are provided. If data tables are not shown, no standing crop estimates were obtained for this lake.




Table 4. Comparison between the average biomass from three zones within your lake and among all lakes assessed within your watershed.

|                 | Lake Helen | Rocky/ Brushy Creek |
|-----------------|------------|---------------------|
|                 |            | (Average)           |
| Emergent Zone:  | 13.92      | 4.74                |
| Floating Zone:  | 0.00       | 0.69                |
| Submersed Zone: | 1.06       | 1.54                |

Number of lakes sampled in your watershed: 29

Note: All biomass measurements are shown in kilograms per square meter.

Table 5. Dominant taxa from three zones within your lake.

| <u>Zone</u>    | Dominant Plant | <u>Status</u> |
|----------------|----------------|---------------|
| Emergent Zone: | Torpedo Grass  | Exotic        |
| Floating Zone: |                |               |

Submersed Zone: Baldwin's Spikerush, Roadgrass Native